
51gure -

$$\begin{array}{c} CH_3O + CH_3O +$$

2. NaBH₄










BEST AVAILABLE COPY

- 217611


BEST AVAILABLE COPY









• 3




| TOTALL | | | | | | | |
|---------|------------|------------|------------|------------|------------|------------|---|
| yrosine | Residue at | |
| Inase | Region A | Region B | Region C | Region D | Region E | Region F | |
| AK3 | Pro906 | Tyr304 | Leu905 | Met902 | Ala966 | Asp912 | - |
| AK2 | Pro933 | Tyr931 | Leu932 | Met929 | Gly993 | Asp939 | |
| IAK1 | Pro948 | Phe948 | Leu947 | Met944 | Gly1008 | Glu954 | |
| × | Ala478 | Tyr 478 | Met 477 | Thr 474 | Ser 538 | Asn 484 | |
| SYK | Glu342 | Met340 | Ala341 | Met338 | Ser403 | Lys348 | |
| X | Ala342 | Phe340 | Met341 | Ala342 | Ala403 | Asp348 | |
| z | Ala323 | Tyr321 | Met322 | Ala323 | Ala384 | Asp329 | |
| ¥ | Ala1080 | Leu1078 | Met1079 | Met1076 | Gly1149 | Ser1086 | |


JEST AVAILABLE COPY










BEST AVAILABLE COPY






BEST AVAILABLE COPY






4


BEST AVAILABLE COPY








BEST AVAILABLE COPY


12,














BEST AVAILABLE COPY




BEST AVAILABLE COPY














Figur -




P131 provides apoptosis in spilen cells






Days post BMT

131,132 1 sourial







Days post BMT


WHI-P131 (60 mg/kg/day) - effects on GVHD development


| Cumulative proportion surviving | Control WHI-P131 | 51 90.2 \pm 4.2 17.6 \pm 5.3 2.0 \pm 1.9 44 | 31 9 100 \pm 0 77.8 \pm 13.9 33.3 \pm 15.7 82 0.0003 | 98.9 ± 10.5 11.1 ±10.5 0 ±0 44 0.485 | exate 16 93.8 \pm 6.1 56.2 \pm 12.4 25.0 \pm 10.8 63.5 0.0009 0.356 | exate + WHI-P131 13 100 \pm 0 76.9 \pm 11.7 61.5 \pm 13.5 87 <0.0001 0.320 | orine A 15 73.3 \pm 11.4 26.7 \pm 11.4 13.3 \pm 8.8 45 0.493 0.029 | orine A \pm WHI-P131 15 80.0 \pm 10.3 33.3 \pm 12.2 20.0 \pm 10.3 51 0.198 0.067 |
|---------------------------------|--------------------|---|--|--|---|--|--|--|
| | Group | CONTROL | WHI-P131 | WHI-P132 | M thotrexate | Methotrexate + WHI-P131 | Cyclosporine A | Cyclosporine A + WHI-P131 |


Methotrexate vs.
Methotrexate + WHI-P131
0.062






















tigure >














Days post transplantation







V11

